

Dependence of the large-scale vortex instability on latitude, stratification, and domain size

M.J. Mantere^{1,*}, P.J. Käpylä^{1,2}, and T. Hackman^{1,3}

¹ Department of Physics, PO BOX 64 (Gustaf Hällströmin katu 2a), FI-00014 University of Helsinki, Finland

² NORDITA, AlbaNova University Center, Roslagstullsbacken 23, SE-10691 Stockholm, Sweden

³ Finnish Centre for Astronomy with ESO, University of Turku, Väisäläntie 20, FI-21500 Piikkiö, Finland

Received 2011 Sep 20, accepted 2011 Nov 7

Published online 2011 Dec 12

Key words convection – hydrodynamics – methods: numerical – turbulence

In an earlier study, we reported on the excitation of large-scale vortices in Cartesian hydrodynamical convection models subject to rapid enough rotation. In that study, the conditions for the onset of the instability were investigated in terms of the Reynolds (Re) and Coriolis (Co) numbers in models located at the stellar North pole. In this study, we extend our investigation to varying domain sizes, increasing stratification, and place the box at different latitudes. The effect of the increasing box size is to increase the sizes of the generated structures, so that the principal vortex always fills roughly half of the computational domain. The instability becomes stronger in the sense that the temperature anomaly and change in the radial velocity are observed to be enhanced. The model with the smallest box size is found to be stable against the instability, suggesting that a sufficient scale separation between the convective eddies and the scale of the domain is required for the instability to work. The instability can be seen up to the colatitude of 30 degrees, above which value the flow becomes dominated by other types of mean flows. The instability can also be seen in a model with larger stratification. Unlike the weakly stratified cases, the temperature anomaly caused by the vortex structures is seen to depend on depth.

© 2011 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim

1 Introduction

Hydrodynamical Cartesian convection simulations subject to high enough rotational influence ($Co \gtrsim 3$) and large enough Reynolds number ($Re \gtrsim 30$) have been reported to generate vortices, the sizes of which are large compared to the size of the convection cells (e.g. Chan 2003, 2007; Käpylä, Mantere & Hackman 2011, hereafter KMH11). In the moderate Coriolis number regime, the vortices are cyclonic, suppressing the energy transport by convection, and thereby appearing as cooler than their surroundings. When Coriolis number is further increased, anticyclonic vortices are preferred, enhancing the convective energy transport, making the vortices appear as regions warmer than their surroundings (KMH11).

In our previous study (KMH11), we proposed that such vortical structures could be responsible for cool/hot starspots in rapidly rotating late-type stars possessing outer convection zones. We were prompted to look into such a possibility due to the decorrelation of the surface temperature maps, obtained by Doppler-imaging techniques, from the distribution of surface magnetic fields, derived through Zeeman-Doppler imaging methods (e.g. Donati & Collier Cameron 1997; Donati 1999; Hussain et al. 2000; Jeffers et al. 2011; Kochukhov et al. 2011).

The resulting temperature anomaly due to the vortices was shown to be of the order of 5%, being somewhat

weaker than the temperature contrasts deduced from observations. The model, however, was very simple: for example, the density stratification in the radial direction was only of the order of 23. Due to the small growth rate of the instability, requiring several thousand turnover times to saturate, only a very limited parameter range was studied. In this study, we extend the previous one by investigating a model with a larger stratification, models with varying domain size, and place the computational domains at different latitudes.

2 Model

Our model is based on that used by Käpylä et al. (2009) and KMH11. A rectangular portion of a star is modeled by a box situated at colatitude θ . The box is divided into three layers: an upper cooling layer, a convectively unstable layer, and a stable overshoot layer (see below). We solve the following set of equations for compressible hydrodynamics:

$$\frac{D \ln \rho}{Dt} = -\nabla \cdot \mathbf{U}, \quad (1)$$

$$\frac{D\mathbf{U}}{Dt} = -\frac{1}{\rho} \nabla p + \mathbf{g} - 2\boldsymbol{\Omega} \times \mathbf{U} + \frac{1}{\rho} \nabla \cdot 2\nu\rho\mathbf{S}, \quad (2)$$

$$\frac{De}{Dt} = -\frac{p}{\rho} \nabla \cdot \mathbf{U} + \frac{1}{\rho} \nabla \cdot K \nabla T + 2\nu\mathbf{S}^2 - \frac{e - e_0}{\tau(z)}, \quad (3)$$

where $D/Dt = \partial/\partial t + \mathbf{U} \cdot \nabla$ is the advective time derivative, ν is the kinematic viscosity, K is the heat conductivity,

* Corresponding author: maarit.mantere@helsinki.fi

ρ is the density, \mathbf{U} is the velocity, $\mathbf{g} = -g\hat{z}$ is the gravitational acceleration, and $\boldsymbol{\Omega} = \Omega_0(-\sin\theta, 0, \cos\theta)$ is the rotation vector. The fluid obeys an ideal gas law $p = (\gamma-1)\rho e$, where p and e are the pressure and the internal energy, respectively, and $\gamma = c_P/c_V = 5/3$ is the ratio of the specific heats at constant pressure and volume, respectively. The specific internal energy per unit mass is related to the temperature via $e = c_V T$. The rate of the strain tensor \mathbf{S} is given by

$$S_{ij} = \frac{1}{2}(U_{i,j} + U_{j,i}) - \frac{1}{3}\delta_{ij}\nabla \cdot \mathbf{U}. \quad (4)$$

The last term of Eq. (3) describes cooling at the top of the domain. Here $\tau(z)$ is a cooling time which has a profile smoothly connecting the upper cooling layer and the convectively unstable layer below, where $\tau \rightarrow \infty$.

The positions of the bottom of the box, bottom and top of the convectively unstable layer, and the top of the box, respectively, are given by $(z_1, z_2, z_3, z_4) = (-0.85, 0, 1, 1.15)d$, where d is the depth of the convectively unstable layer. In the case of larger stratification (Set C), the corresponding vertical positions are $(z_1, z_2, z_3, z_4) = (-0.4, 0, 1, 1.1)d$, resulting in a vertical extent somewhat smaller than in Sets A and D. Initially the stratification is piecewise polytropic with polytropic indices $(m_1, m_2, m_3) = (3, 1, 1)$, which leads to a convectively unstable layer above a stable layer at the bottom of the domain. In a system set up this way, convection transports roughly 20 per cent of the total flux. Due to the presence of the cooling term, a stably stratified isothermal layer forms at the top. The standard horizontal extent of the box, $L_H \equiv L_x = L_y$, is $4d$; the horizontal domain size is varied from half of this to double the size in Set A. In Set C with larger stratification, the horizontal extent of the box is $5d$. The simulations in Sets A and C are made at the North pole, corresponding to $\theta = 0^\circ$, while in Set D, the colatitude is varied in steps of 15° to cover the latitude range down to $\theta = 60^\circ$. The simulations were performed with the PENCIL CODE¹, which is a high-order finite difference method for solving the compressible equations of magnetohydrodynamics.

2.1 Units and non-dimensional parameters

Non-dimensional quantities are obtained by setting

$$d = g = \rho_0 = c_P = 1, \quad (5)$$

where ρ_0 is the initial density at z_2 . The units of length, time, velocity, density, and entropy are

$$\begin{aligned} [x] &= d, & [t] &= \sqrt{d/g}, & [U] &= \sqrt{dg}, \\ [\rho] &= \rho_0, & [s] &= c_P. \end{aligned} \quad (6)$$

We define the Prandtl number and the Rayleigh number as

$$\text{Pr} = \frac{\nu}{\chi_0}, \quad \text{Ra} = \frac{gd^4}{\nu\chi_0} \left(-\frac{1}{c_P} \frac{ds}{dz} \right)_0, \quad (7)$$

where $\chi_0 = K/(\rho_m c_P)$ is the thermal diffusivity, and ρ_m is the density in the middle of the unstable layer, $z_m =$

$\frac{1}{2}(z_3 - z_2)$. The entropy gradient, measured at z_m , in the non-convective hydrostatic state, is given by

$$\left(-\frac{1}{c_P} \frac{ds}{dz} \right)_0 = \frac{\nabla - \nabla_{\text{ad}}}{H_P}, \quad (8)$$

where $\nabla - \nabla_{\text{ad}}$ is the superadiabatic temperature gradient with $\nabla_{\text{ad}} = 1 - 1/\gamma$, $\nabla = (\partial \ln T / \partial \ln p)_{z_m}$, and where H_P is the pressure scale height. The amount of stratification is determined by the parameter $\xi_0 = (\gamma - 1)e_0/(gd)$, which is the pressure scale height at the top of the domain normalized by the depth of the unstable layer. We use $\xi_0 = 1/3$ in Sets A and B, which results in a density contrast of about 23 across the whole domain, and roughly 9 over the convectively unstable layer. We make one run with higher stratification (C1), for which $\xi_0 = 1/6$, resulting in a density contrast of roughly 230 over the convectively unstable layer.

We define the Reynolds and Péclet numbers via

$$\text{Re} = \frac{u_{\text{rms}}}{\nu k_f}, \quad \text{Pe} = \frac{u_{\text{rms}}}{\chi_0 k_f} = \text{Pr} \text{Re}, \quad (9)$$

where $k_f = 2\pi/d$ is adopted as an estimate for the wavenumber of the energy-carrying eddies, and $u_{\text{rms}} = \sqrt{3u_z^2}$. This definition of u_{rms} neglects the contributions from the large-scale vortices that are generated in the rapid rotation regime (see KMH11). Note that with our definitions, Re and Pe are smaller than the usual ones by a factor of 2π . The amount of rotation is quantified by the Coriolis number, defined as

$$\text{Co} = \frac{2\Omega_0}{u_{\text{rms}} k_f}. \quad (10)$$

We also quote the value of the Taylor number,

$$\text{Ta} = (2\Omega_0 d^2 / \nu)^2, \quad (11)$$

which is related to the Ekman number via $\text{Ek} = \text{Ta}^{-1/2}$.

2.2 Boundary conditions

The horizontal boundaries are periodic for all variables. Stress-free conditions are used for the velocity at the vertical boundaries

$$U_{x,z} = U_{y,z} = U_z = 0. \quad (12)$$

The temperature is kept constant on the upper boundary, and the temperature gradient

$$\frac{dT}{dz} = \frac{-g}{c_V(\gamma-1)(m+1)} \quad (13)$$

is held constant at the lower boundary, yielding a constant heat flux $F_0 = -K\partial T/\partial z$ through the lower boundary.

3 Results

In an earlier study, we investigated the excitation of large-scale vortices in Cartesian domains with weak density stratification, located at the North pole of a rapidly rotating star (KMH11). In that study, we investigated the limiting Reynolds and Coriolis numbers (i.e. inverse Rossby

¹ <http://code.google.com/p/pencil-code/>

Table 1 Summary of the runs. Stars indicate that the simulation has not been run to a saturated state. For Run C1, the kinematic viscosity had to be increased during the last 500 turnover times of the computation to achieve saturation. In this table, however, we list numbers based on the lower value of kinematic viscosity used during the first 2000 turnover times of the simulation. The amount of stratification, $\Delta\rho$, is measured over the convectively unstable region. The temperature anomaly, T_{cyc}/\bar{T} , where T_{cyc} is the extremum of temperature in a cyclonic region and \bar{T} the mean temperature of certain horizontal layer, is measured at the middle of the convective layer, z_m . For runs in which both anti- and cyclonic vortices are present, we list the maximal temperature anomaly caused by either of the structures. The dimensionless input heat flux at the lower boundary of the box is given by $\tilde{F}_0 = F_0/(\rho c_s^2)$, where c_s is the adiabatic sound speed and ρ is the density, both measured at the lower boundary of the domain. The last column indicates the presence of cyclonic (C), anti-cyclonic (A), or both types (A+C) of vortices.

Run	Grid	L_H	θ	$\Delta\rho$	T_{cyc}/\bar{T}	Re	Pe	Pr	Ra	Co	Ta	\tilde{F}_0	Cyc.
D1	$256^2 \times 128$	1	60	9	–	34	16	0.48	2.0×10^6	7.5	1.0×10^8	1.7×10^{-5}	no
D2	$256^2 \times 128$	1	45	9	–	32	16	0.48	2.0×10^6	7.8	1.0×10^8	1.7×10^{-5}	no
D3	$256^2 \times 128$	1	30	9	+0.043	34	16	0.48	2.0×10^6	7.6	1.0×10^8	1.7×10^{-5}	yes (A+C)
D4	$256^2 \times 128$	1	15	9	–0.059	33	16	0.48	2.0×10^6	7.7	1.0×10^8	1.7×10^{-5}	yes (A+C)
D5	$256^2 \times 128$	1	0	9	–0.071	41	19	0.48	2.0×10^6	6.3	1.0×10^8	1.7×10^{-5}	yes (C)
A1*	$512^2 \times 128$	2	0	9	(+0.098)	44	16	0.36	$2.0 \cdot 10^6$	7.7	1.8×10^8	1.7×10^{-5}	yes (2×A)
A2*	$256^2 \times 128$	1	0	9	(+0.062)	42	15	0.36	2.0×10^6	8.1	1.8×10^8	1.7×10^{-5}	yes (A)
A3	$128^2 \times 128$	0.5	0	9	–	39	14	0.36	2.0×10^6	8.7	1.8×10^8	1.7×10^{-5}	no
C1	$256^2 \times 192$	1	0	233	+0.052	94	48	0.5	1.3×10^7	4.3	2.6×10^8	3.4×10^{-5}	yes (A+C)

numbers) above which the instability was excited. For the present study, we have extended our analysis to models with varying computational domain size (Sect. 3.1), place the box at different latitudes of the star (Sect. 3.2), and present a model with higher density stratification (Sect. 3.3). The most relevant input parameters for the runs and some essential output quantities are listed in Table 1. As is evident from this table, some of the runs generating vortices have not yet reached a completely saturated state (these runs are marked with stars), even though they have been integrated for several thousands of turnover times. In such cases, quantities such as the temperature anomaly between the vortex and its surroundings, are underestimated, and the listed numbers in Table 1 should be regarded as lower limits.

3.1 Dependence on the domain size

In our previous computations with the standard box size of $L_H = 4d$ (KMH11), we observed a clear tendency of the sizes of the vortices to approach the wavenumber $k/k_1 = 1$, i.e. they tended to fill in as large a fraction of the horizontal extent as possible. This prompted us to study the dependence of the instability on the horizontal extent of our Cartesian box. In Run A3 of this paper (see Table 1), both the horizontal extent and resolution of the domain are halved. Run A2 is a model with the standard box size (otherwise identical to D5 except for the values of the kinematic viscosity and rotation rate), and in Run A3 the resolution and horizontal extent are doubled. In all the runs, the computational domain is located at the North pole of the star.

In Run A3, the Reynolds (roughly 39) and Coriolis numbers (roughly 8.7) are clearly above the critical values found in KMH11; still, no vortices are excited. This is evident from the leftmost panels of Fig. 1, where we show the tem-

perature field (upper panel) and radial velocity (lower panel) at the middle of the convectively unstable layer, z_m . Some large-scale $k/k_1 = 1$ pattern can be detected in the temperature field, that might be indicative of the early stages of the vortex-instability. This run, however, was continued up to 6500 turnover times, matching the timepoint of the slice plotted in Fig. 1. It is not completely ruled out that a very slowly growing vortex instability mode is present, but in any case its growth rate is strongly reduced from the standard box runs presented in KMH11.

Run A2 shows very similar behavior to the earlier calculations presented in KMH11: the Reynolds number is clearly supercritical to the instability (roughly 42), and Coriolis number (roughly 8.1) in the regime where the excitation of an anti-cyclone, i.e. a vortex rotating in opposite direction to the overall rotation of the domain, was reported. As shown in the middle panels of Fig. 1, a vortex rotating in the clock-wise direction is seen in the velocity field (lower panel), the structure being warmer than its surroundings (upper panel). The temperature anomaly is slightly more than six percent, a value slightly larger to the number reported in KMH11. We note that, even though the run has been continued up to 5400 turnover times, the run has not yet completely saturated. The temperature across the vortex at z_m , normalised to the mean temperature of the horizontal layer, is plotted in the upper panel of Fig. 2 with a red, dashed line. In the region of the anti-cyclone, the vertical velocities become somewhat enhanced, as evident from the lower panel of the same figure, where the velocity profile normalised to the rms velocity of the horizontal layer across the vortex is plotted with a red, dashed line.

In the model with the largest horizontal extent, Run A1, exhibiting similar Reynolds and Coriolis numbers in comparison to the standard box Run A2, two instead of one an-




Fig. 1 (online colour at: www.an-journal.org) Temperature (*upper row*) and radial velocity U_r (*lower row*) in the middle of the convectively unstable layer, i.e. at z_m , for Runs A3, A2 and A1 (*from left to right*). For Run A3, the slice is taken from the time $t = 6400 \tau_{to}$, where $\tau_{to} = (u_{rms} k_f)^{-1}$. For Runs A2 and A1, the snapshots are taken at $3400 \tau_{to}$ and $2700 \tau_{to}$.

tyclone are seen. The growth rate of the instability, measured from the growth of the horizontal velocity components, increases by a factor of 2.3 from Run A2 to A1. As evident from the rightmost panels of Fig. 1, the two structures are of unequal strength, possibly suggesting that the system has not yet completely saturated, although it has been followed for up to nearly 3700 turnover times. Based on earlier experience with the standard box runs (KMH11), where multiple vortices were also seen, when followed long enough, only one vortex very close to the $k/k_1 = 1$ wavenumber would be the stable end point. The temperature and vertical velocity anomaly across the stronger vortex is plotted in Fig. 2 with solid, black lines. As can be seen from this figure, the temperature anomaly in the middle of the vortex is roughly twice as large as in the standard box case, suggesting that the strength of the instability is indeed dependent on the box size. No such dramatic difference can be seen in the vertical velocity cut across the vortex (Fig. 2 lower panel).

The size of the stronger vortex in Run A1, again, is nearly half of the domain size, i.e. approaching the $k/k_1 = 1$ mode. In Fig. 3 we plot the power spectra for the kinetic energy from Runs A1 (black lines) and A2 (red lines); the wavenumber scale for Run A1 is re-scaled to match

the one of Run A2. In the early stages, when no vortices are yet excited (dashed linestyles), both the power spectra consistently peak at intermediate wavenumber of roughly $k/k_1 = 7$; this number reflects the size of the turbulent eddies due to convective motions. Due to the appearance of the vortices, the energy contained in large scales grows, and dominates the flow in the nearly saturated stage. Most of the power is seen near the wavenumber $k/k_1 = 1$. In Run A1, the instability would still have 'space' to advance into even lower wavenumbers i.e. larger scales; followed up even further, it might still do so.

3.2 Latitudinal dependence

In Set D, we place the computational domain at different colatitudes with a coarse latitude grid of $[0, 15, 30, 45, 60]$ using the standard box size. In all the runs in Set D, the Reynolds and Coriolis numbers are kept above the critical values found by KMH11. We note that the Reynolds numbers realized in the runs are quite close to the critical values of KMH11, but due to numerical stability, it was impossible to use lower values of kinematic viscosity without the cost of having to increase spatial resolution as well. This, on the other hand, was not possible due to the prohibitive




Fig. 2 (online colour at: www.an-journal.org) Temperature (*upper panel*) and vertical velocity U_r (*lower panel*) across the anticyclone for Runs A1 and A2 with differing box size. The cuts are taken from the depth z_m , i.e. the middle of the convection zone.




Fig. 3 (online colour at: www.an-journal.org) Kinetic energy spectra for the Runs A1 (black lines) and A2 (red lines). Spectra taken at early times, when no vortices are yet excited, are plotted with dashed linestyle, and spectra from the vortex-state with solid linestyle. The wavenumber range of Run A1 is scaled to match the one of Run A2.




Fig. 4 (online colour at: www.an-journal.org) Two-dimensional slice, in the xz -plane, of the azimuthal velocity U_y from the run D3 at colatitude 30 degrees. The anticyclone generated in the run shows up as a structure spanning through the entire vertical extent of the computational domain, inclined with the rotation vector.

computational cost of the runs even at present resolutions. In Runs D5, D4 and D3, we still observe the excitation of vortices, but in the rest, other types of large-scale flows are generated, suppressing the instability. Such large-scale flows are normally referred to as banana cells, seen both in Cartesian (Chan 2001; Käpylä et al. 2004) and spherical geometries (e.g. Brown et al. 2008; Käpylä et al. 2011a, 2011b).

The structures extend through the whole convection zone and even penetrate to the overshoot region, see Fig. 4, where we plot a radial-vertical (xz) slice of the azimuthal velocity U_y from Run D3 with $\theta = 30^\circ$. The convection, and also the vortex tube, are strongly affected by rotation, which forces the structures to become inclined with the axis of rotation.

The growth rate of the instability is reduced when the colatitude is increased; therefore, it has been extremely difficult to follow the Runs D3 and D4 up to saturation. All of the runs in Set D, however, have been run up to near saturation, when the growth of the horizontal velocity components slows down. The run at the pole (D5) having the largest growth rate is the only run in Set D showing only one single type of structure, namely a cyclone. In an earlier state, this run also exhibited both an anti-cyclonic and cyclonic vortex; in the later stages, however, only the cyclone persists. The other runs still show both types of vortices, which might still be indicative of the runs being further away from complete saturation than D5. It is also noteworthy, that in addition of the growth rate getting smaller as function of colatitude, the temperature anomaly shows a similar trend, i.e. the temperature contrast is getting weaker the further away one moves from the pole.

3.3 Dependence on stratification

We have made an attempt to quantify the effect of increasing stratification on the vortex instability by running one model

where the stratification is almost 30 times larger than in our standard cases. Due to this, larger resolution in the radial direction is needed; the amount of gridpoints has been increased from $n_z = 128$ to 192, making these computations even more demanding than the rest included in this study. This run is identified as Run C1 in Table 1, and can be observed to have a much higher Reynolds number (roughly 94) than any other run, but clearly a lower Coriolis number (roughly 4.3) than the rest of the runs. Nevertheless, these numbers exceed the critical values found in KMH11, and therefore the vortex-instability is to be expected, unless the increased stratification has a significant effect on its excitation conditions.

Indeed, we observe the instability in Run C1, generating a stronger cyclonic and a weaker anti-cyclonic vortex, although the growth rate of it is reduced in comparison to the other runs due to the lower Coriolis number. We have been able to calculate this model up to roughly 2000 turnover times, but it is clear that the system is still far from saturation. With the value of kinematic viscosity used, the system becomes numerically unstable quite soon after this timepoint; increasing the viscosity twofold, the system is observed to saturate after about 2500 turnover times. The main properties, such as the size and temperature contrast of the vortices, do not considerably change as the viscosity is increased. In Table 1 we list numbers with the lower viscosity coefficient for the first 2000 turnover times of the simulation.

The temperature anomaly measured before saturation, using the lower viscosity coefficient, and at the saturated state, achieved by increasing the kinematic viscosity, is close to 5%. This number is comparable to the standard box calculations very near saturation, and the values obtained in KMH11. The vortices, again, occur at the very largest scales of the box, and their size does not vary significantly as a function of depth, even though the system is more strongly stratified. The temperature anomaly, on the contrary, varies monotonically through the convection zone, see Fig. 5, where we plot a temperature cut in the y -direction through a cyclonic, cooler, region located at $x = -1.1d$ for various depths. From this figure it is evident that the temperature contrast is monotonically increasing with height. In the models with weaker stratification, such an effect is not clearly visible.

4 Conclusions

In an earlier study (KMH11) we reported on the excitation of large-scale vortices in Cartesian hydrodynamical convection models subject to rapid enough rotation. In that study, the conditions of the onset of the instability were investigated in terms of the Reynolds and Coriolis numbers in models located at the stellar North pole. In this study, we extend our investigation to varying domain sizes, increasing stratification and place the box at different latitudes.




Fig. 5 (online colour at: www.an-journal.org) Azimuthal cuts through the computational domain at $x = -1.1d$ for various depths for the Run C1. The bluer the color, the nearer the bottom of the convection zone the cut is taken. Red colors represent cuts near z_m , and the black line a cut taken exactly at z_m . The yellow colors represent cuts taken near the top of the convection zone.

The effect of the increasing box size is to increase the sizes of the generated structures, so that the principal vortex always fills roughly half of the computational domain. The instability becomes stronger in the sense that the temperature anomaly and change in the radial velocity are observed to be enhanced. Also the growth rate, measured from the time evolution of the horizontal velocity components, is more than doubled when the boxsize is doubled. The model with the smallest box size is found to be stable against the instability even though the critical Reynolds and Coriolis numbers found in the earlier study KMH11 are clearly exceeded, suggesting that a sufficient scale separation between the convective eddies and the smallest wavenumber of the domain is required for the instability to work.

The instability can be seen up to a colatitude of 30 degrees, but at higher colatitudes, the flow becomes dominated by large-scale flows known as banana cells. Such flows have earlier been found from Cartesian and spherical convection models. The vortices are seen to align with the rotation vector, forming structures tilted but coherent through the entire convection zone, extending even to the overshoot region. The growth rate of the instability is observed to become lower with increasing colatitude. It also seems evident that the temperature contrast of the vortices is getting weaker the further away one moves from the pole.

Only very little variance of the temperature contrast across the vortices can be seen as function of height when stratification is small. The instability can also be seen in a model with larger stratification. Unlike the weakly stratified cases, the temperature anomaly caused by the vortex structures is seen to depend on height.

Acknowledgements. Computational resources granted by UCSC – IT Center for Science, financed by the Ministry of Education, and financial support from the Academy of Finland grants No. 136189,

140970 (PJK) and 218159, 141017 (MJM), and the 'Active Suns' research project at University of Helsinki (TH) is acknowledged. The authors acknowledge the hospitality of NORDITA during their visits.

References

- Brown, B.P., Browning, M.K., Brun, A.S., Miesch, M.S., Toomre, J.: 2008, *ApJ* 689, 1354
- Chan, K.L.: 2001, *ApJ* 548, 1102
- Chan, K.L.: 2003, in: S. Turcotte, S.C. Keller, R.M. Cavallo (eds.), *3D Stellar Evolution*, ASPC 293, p. 168
- Chan, K.L.: 2007, *AN* 328, 1059
- Donati, J.-F.: 1999, *MNRAS* 302, 457
- Donati, J.-F., Collier Cameron, A.: 1997, *MNRAS* 291, 1
- Hussain, G.A.J., Donati, J.-F., Collier Cameron, A., Barnes, J.R.: 2000, *MNRAS* 318, 961
- Jeffers, S.V., Donati, J.-F., Alecian, E., Marsden, S.C.: 2011, *MNRAS* 411, 1301
- Käpylä, P.J., Korpi, M.J., Tuominen, I.: 2004, *A&A* 422, 793
- Käpylä, P.J., Korpi, M.J., Brandenburg, A.: 2009, *ApJ* 697, 1153
- Käpylä, P.J., Mantere, M.J., Hackman, T.: 2011, *ApJ* 742, 34 (KMH11)
- Käpylä, P.J., Korpi, M.J., Guerrero, G., Brandenburg, A., Chatterjee, P.: 2011a, *A&A* 531, A162
- Käpylä, P.J., Mantere, M.J., Brandenburg, A.: 2011b, *AN* 332, 883
- Kochukhov, O., Hackman, T., Mantere, M.J., Ilyin, I., Piskunov, N., Tuominen, I.: 2011, in preparation